Beginners' Guide to Macro Moths Te Hiku

















Te Takenga mai o te Pepe -

Tane climbs to the highest

heaven to visit his brother, Rehua.

Rehua calls for a fire to be lit and

Tāne asks his brother "What will we

be eating, brother?" At this, Rehua

calabashes to be brought forth.

shakes his head and two koko/

tūi fall from his hair. Tāne refuses

to eat the birds as they have been

living in Rehua's hair and are tapu.

Tane asks if he can return to earth

with the birds. Rehua agrees, and

says that the birds will live in the

trees, and feed on the fruits of the

trees. When Rehua shook his head

other 'birds' also fell out: pepee,

pihareinga, the grasshopper; and

kēkerewai, the beetle. Collectively

these are known as 'Ngā Manu a

Rehua' and come out in summer.

the moth; tātarakihi, the cicada;

The Origin of the Moth

Ngā Whetū - The Stars

The story of the moth is told in the stars of the Rehua constellation. Rehua, the eldest son of Raki and Papatūānuku, is the reddish star Antares in the western Scorpius constellation.

Wairua Tangata -

A common held belief was that the spirits of those who had passed would be reincarnated as a moth.

Moths are an important part of New Zealand's biodiversity

Lepidoptera (moths and butterflies) are the third largest group of insects in New Zealand (over 1750 named species). Scientists estimate there are still more than 300 moth species. to be discovered and named in New Zealand.

'The Spirit of a Person'

Ka hou ki te whenua, he tūkoukou; ka puta ki te rangi he pepe -The pupa enters the Earth, a moth emerges to the sky

NZ moths are special

them.

More than 86% of the known moths in NZ are endemic. They only occur in NZ so we have to look after

Why are moths important?

Moths are a key part of the wider ecosystem and they sit in the centre of a complex food web. The caterpillars are herbivores eating a range of native plants.

Many adult moths drink nectar and are important pollinators. Many of our native plants are likely pollinated by moths. Moths go quietly about this critical job during the night when most people are sleeping, which means we often overlook how important they are. Moths are a major food source for other insects and native birds. Unfortunately, they are also an important food source for introduced pests like mice, rats, and

Moths can also be pests

hedgehogs.

Hīhue (the kūmara moth, Agrius convolvuli) had a large impact on kūmara crops and there are

many whakatauki written about the caterpillar and its capacity to eat.

A more recent pest is the codling moth (Cydia pomonella). Introduced from Europe, it attacks apples, pears, walnuts, and other fruit.

Moths have a bad reputation for eating clothes, especially natural fibres like wool, silk, and fur. In reality there are very few moth species whose caterpillars eat clothes. You can easily solve this problem by putting the clothes in and out of the freezer a couple of times.

Puka Whakamārama o te Pepe Nui This guide contains only the most common larger moths in this large region. If you find a moth that is not in this guide, this may be because it is rare, a range extension (not normally found in this place), a 'micro-moth', an introduced species or a new species. Please let us know.

Who benefits from studying moths? Educatio Stories Knowledge Biosecurity Appreciatio Music Kaitiakita Pollination

We can use science to understand ecosystem responses to land-use change, climate change and light pollution We need to know what we have so we can better protect it

Science

Ecology

Biogeography

Taxonom

The more we know about things, the more we appreciate their beauty and intrinsic value

Ecosyster

ight for eel

most people don't see how many there are or what's happening to them. Unfortunately we need specimens for identification.

Why we need standardised data

If we all use the same type of trap (e.g. a Heath Moth Trap) we can compare data from different places and over time. With standardised information we can all work together to keep an eye on the moths. This way, if they start to decline we can see something is happening and work together to take better care of the moths.

Why we need quantitative data Counting the number of individuals of each species gives us more information than just a list of the species present. This quantitative information helps to detect change over time or between places.

Vhen the pepetuna are in great numbers it is a good

Te āwheto kai paeka – The caterpillar eats the leaf (Said of someone who goes around tasting various dishes) Me te upoko tūkoukou - like the chrysalis of the Sphinx moth (When likening something to especially fine weaving)

I hikaina te ahi titi, engari i whakapoapoa kē i te pepe -The fire was lit to attract the mutton birds and the moths flew into it – Te Whiti o Rongomai

Why we need to trap the moths Moths are mainly out at night, so

Why do we need to study moths?

Moths breed fast and have lots of offspring. There are lots of different moth species playing different roles in the ecosystem. Moths have links to lots of other species (e.g. plants, birds, introduced pests, other invertebrates).

If something is changing in the ecosystem, moths are amongst the first creatures to respond. They are likely to be good indicators of change.

What does that word mean?

Lepidoptera: moths and butterflies Endemic: found only in that place *Ecosystem*: all biological and physical processes in an area Food web: what eats what Herbivores: animals that eat plants *Nocturnal*: active at night *Community*: species in an area



Where to get more information & help www.landcareresearch.co.nz/mothnet Email: MothNetOtago@gmail.com

@MothNetNZ

f MothNetNZ

www.naturewatch.org.nz/ Entomological Society of New Zealand: www.ento.org.nz

Acknowledgements

The text, content, and design are by the Ahi Pepe MothNet project partners: Dr Barbara J Anderson, Komene Cassidy, Victoria Campbell, Dr Robert Hoare, Tiahuia Kawe-Small, Tahu MacKenzie, Dr Ralf Ohlemüller, Tangiwai Rewi, Dr Priscilla Wehi and Emma Burns. Illustrations are by Sean W Gilles. The moth list was compiled by Dr Robert Hoare, with help from NZ's lepidopterist community. The moth images are photographs by Birgit Rhode of specimens in the NZAC, taken with funding from the Terrestrial Freshwater Biodiversity Information System (TFBIS) programme.

This guide was produced by Manaaki Whenua -Landcare Research as part of the Ahi Pepe MothNet project funded by the MBIE Unlocking Curious Minds initiative and Biological Heritage National Science Challenge.



URIOUS



